

Estimated chronic wasting disease (CWD) prevalence¹ from harvested male deer. (See notes at right for further explanation.)

Game Management Unit (GMU)	GMU-level estimated prevalence ² (from mandatory submissions)	Prevalence Range ³	Herd/DAU
001	[5%]	5-10%	D-01
002	[5%]	5-10%	D-01
003	11%	10-20%	D-02
004	31%	over 20%	D-02
005	[21%]	over 20%	D-02
006	[15%]	10-20%	D-03
007	[5%]	under 5%	D-04
008	[5%]	under 5%	D-04
009	[5%]	under 5%	D-04
010	[4%]	under 5%	D-06
011	19%	10-20%	D-07
012	36%	over 20%	D-07
013	[24%]	over 20%	D-07
014	[18%]	10-20%	D-02
015	6%	5-10%	D-08
016	[15%]	10-20%	D-03
017	[15%]	10-20%	D-03
018	[8%]	5-10%	D-09
019	8%	5-10%	D-04
020	8%	5-10%	D-10
021	3%	under 5%	D-11
022	9%	5-10%	D-07
023	22%	over 20%	D-07
024	[24%]	over 20%	D-07
025	[7%]	5-10%	D-43
026	[7%]	5-10%	D-43
027	[8%]	5-10%	D-09
028	[8%]	5-10%	D-09
029	[19%]	10-20%	D-27
030	[3%]	under 5%	D-11
031	7%	5-10%	D-41
032	[7%]	5-10%	D-41
033	9%	5-10%	D-42
034	[7%]	5-10%	D-43
035	3%	under 5%	D-08
036	2%	under 5%	D-08
037	[8%]	5-10%	D-09
038	[19%]	10-20%	D-27
039	[3%]	under 5%	D-17
040	1%	under 5%	D-18
041	[5%]	5-10%	D-12
042	8%	5-10%	D-12
043	[3%]	under 5%	D-13
044	<1%	not detected	D-14

Notes

¹ Prevalence means the proportion of sampled animals infected by CWD calculated as [number infected / total number sampled] and expressed here as a percentage (e.g., 10%). A prevalence of 20% is the same as a ratio of 1 in 5 (i.e. 1 in every 5 animals tested is infected), 10% is 1 in 10, 5% is 1 in 20, 1% is 1 in 100.

² CPW's best estimate of prevalence based on testing 100 or more submissions from this Game Management Unit (GMU). Values in brackets [] are based on the overall herd (Data Analysis Unit) prevalence estimate. Estimates shown for mule deer (MD) and white-tailed deer (WTD) where different. Data derived from the most recent year(s) when sample submissions were required in the GMU.

³ Range of estimated prevalence within the corresponding herd/Data Analysis Unit (DAU) based on at least one year of mandatory sample submissions regardless of sample size. **Bold (5-10%)** or **bold red (10-20%;shaded >20%)** denotes GMUs/DAUs that exceed the 5% threshold that now compels CWD management action under CPW's response plan.

"Not detected" means no cases reported from this DAU (herd) as of January 2023. **Does not assure absence of CWD.**

Estimated chronic wasting disease (CWD) prevalence¹ from harvested male deer. (See notes at right for further explanation.)

Game Management Unit (GMU)	GMU-level estimated prevalence ² (from mandatory submissions)	Prevalence Range ³	Herd/DAU	Notes
045	[3%]	under 5%	D-08	
046	[3%]	under 5%	D-17	
047	[3%]	under 5%	D-13	
048	not detected	not detected	D-15	
049	[<1%]	under 5%	D-16	
050	[1%]	under 5%	D-38	
051	<1%	under 5%	D-17	
052	10%	10-20%	D-51	
053	not detected	not detected	D-20	
054	not detected	not detected	D-21	
055	not detected	not detected	D-22	
056	not detected	not detected	D-15	
057	[<1%]	under 5%	D-16	
058	[<1%]	under 5%	D-16	
059	[5%]	5-10%	D-50	
060	27%	over 20%	D-23	
061	[14%]	10-20%	D-19	
062	15%	10-20%	D-19	
063	not detected	not detected	D-20	
064	[4%]	under 5%	D-40	
065	2%	under 5%	D-40	
066	not detected	not detected	D-25	
067	not detected	not detected	D-25	
068	not detected	not detected	D-26	
069	<1%	under 5%	D-34	
070	3%	under 5%	D-24	
071	[3%]	under 5%	D-24	
072	[2%]	under 5%	D-29	
073	[2%]	under 5%	D-29	
074	not detected	not detected	D-52	
075	[<1%]	under 5%	D-30	
076	not detected	not detected	D-36	
077	[<1%]	under 5%	D-30	
078	<1%	under 5%	D-30	
079	not detected	not detected	D-36	
080	not detected	not detected	D-35	
081	not detected	not detected	D-35	
082	not detected	not detected	D-37	
083	not detected	not detected	D-31	
084	<1%	under 5%	D-34	
085	1%	under 5%	D-32	
086	[<1%]	under 5%	D-34	
087	[24%]	over 20%	D-05	
088	[24%]	over 20%	D-05	

¹Prevalence means the proportion of sampled animals infected by CWD calculated as [number infected / total number sampled] and expressed here as a percentage (e.g., 10%). A prevalence of 20% is the same as a ratio of 1 in 5 (i.e. 1 in every 5 animals tested is infected), 10% is 1 in 10, 5% is 1 in 20, 1% is 1 in 100.

²CPW's best estimate of prevalence based on testing 100 or more submissions from this Game Management Unit (GMU). Values in brackets [] are based on the overall herd (Data Analysis Unit) prevalence estimate. Estimates shown for mule deer (MD) and white-tailed deer (WTD) where different. Data derived from the most recent year(s) when sample submissions were required in the GMU.

³Range of estimated prevalence within the corresponding herd/Data Analysis Unit (DAU) based on at least one year of mandatory sample submissions regardless of sample size. **Bold (5-10%)** or **bold red (10-20%;shaded >20%)** denotes GMUs/DAUs that exceed the 5% threshold that now compels CWD management action under CPW's response plan.

⁴"Not detected" means no cases reported from this DAU (herd) as of January 2023. **Does not assure absence of CWD.**

Estimated chronic wasting disease (CWD) prevalence¹ from harvested male deer. (See notes at right for further explanation.)

Game Management Unit (GMU)	GMU-level estimated prevalence ² (from mandatory submissions)	Prevalence Range ³	Herd/DAU
089	[24%]	over 20%	D-05
090	[24%]	over 20%	D-05
091	[26%]	over 20%	D-44
092	[26%]	over 20%	D-44
093	[37%]	over 20%	D-54
094	[26%]	over 20%	D-44
095	[24%]	over 20%	D-05
096	[26%]	over 20%	D-44
097	[37%]	over 20%	D-54
098	[37%]	over 20%	D-54
099	[37%]	over 20%	D-54
100	[37%]	over 20%	D-54
101	[40%]	over 20%	D-55
102	[40%]	over 20%	D-55
103	[27%]	over 20%	D-47
104	[11%]	10-20%	D-49
105	9%	5-10%	D-49
106	[11%]	10-20%	D-49
107	[16%]	10-20%	D-46
109	[27%]	over 20%	D-47
110	[6%]	5-10%	D-48
111	[6%]	5-10%	D-48
112	[16%]	10-20%	D-46
113	[16%]	10-20%	D-46
114	[16%]	10-20%	D-46
115	[16%]	10-20%	D-46
116	[27%]	over 20%	D-47
117	[27%]	over 20%	D-47
118	[6%]	5-10%	D-48
119	[6%]	5-10%	D-48
120	[16%]	10-20%	D-46
121	[16%]	10-20%	D-46
122	[8%]	5-10%	D-28
123	[6%]	5-10%	D-48
124	[6%]	5-10%	D-48
125	[8%]	5-10%	D-28
126	[8%]	5-10%	D-28
127	[8%]	5-10%	D-28
128	<1%	under 5%	D-45
129	<1%	under 5%	D-45
130	[8%]	5-10%	D-28
131	[24%]	over 20%	D-07
132	[8%]	5-10%	D-28
133	<1%	under 5%	D-45

Notes

¹ Prevalence means the proportion of sampled animals infected by CWD calculated as [number infected / total number sampled] and expressed here as a percentage (e.g., 10%). A prevalence of 20% is the same as a ratio of 1 in 5 (i.e. 1 in every 5 animals tested is infected), 10% is 1 in 10, 5% is 1 in 20, 1% is 1 in 100.

² CPW's best estimate of prevalence based on testing 100 or more submissions from this Game Management Unit (GMU). Values in brackets [] are based on the overall herd (Data Analysis Unit) prevalence estimate. Estimates shown for mule deer (MD) and white-tailed deer (WTD) where different. Data derived from the most recent year(s) when sample submissions were required in the GMU.

³ Range of estimated prevalence within the corresponding herd/Data Analysis Unit (DAU) based on at least one year of mandatory sample submissions regardless of sample size. **Bold (5-10%) or bold red (10-20%;shaded >20%)** denotes GMUs/DAUs that exceed the 5% threshold that now compels CWD management action under CPW's response plan.

"Not detected" means no cases reported from this DAU (herd) as of January 2023. **Does not assure absence of CWD.**

Estimated chronic wasting disease (CWD) prevalence¹ from harvested male deer. (See notes at right for further explanation.)

Game Management Unit (GMU)	GMU-level estimated prevalence ² (from mandatory submissions)	Prevalence Range ³	Herd/DAU	Notes
134	<1%	under 5%	D-45	
135	<1%	under 5%	D-45	
136	<1%	under 5%	D-45	
137	[8%]	5-10%	D-28	
138	[8%]	5-10%	D-28	
139	[8%]	5-10%	D-28	
140	[1%]	under 5%	D-32	
141	<1%	under 5%	D-45	
142	<1%	under 5%	D-45	
143	not detected	not detected	D-33	
144	not detected	not detected	D-33	
145	not detected	not detected	D-33	
146	[8%]	5-10%	D-28	
147	<1%	under 5%	D-45	
161	[15%]	10-20%	D-03	
171	[15%]	10-20%	D-03	
181	[8%]	5-10%	D-09	
191	3%	under 5%	D-04	
201	[5%]	5-10%	D-01	
211	30%	over 20%	D-07	
214	[21%]	over 20%	D-02	
231	[24%]	over 20%	D-07	
301	[21%]	over 20%	D-02	
361	[3%]	under 5%	D-08	
371	[8%]	5-10%	D-09	
391	[3%]	under 5%	D-17	
411	[7%]	5-10%	D-51	
421	3%	under 5%	D-12	
441	[21%]	over 20%	D-02	
444	<1%	under 5%	D-53	
461	[3%]	under 5%	D-17	
471	[3%]	under 5%	D-13	
481	not detected	not detected	D-15	
500	[1%]	under 5%	D-38	
501	1%	under 5%	D-38	
511	[5%]	5-10%	D-50	
512	[5%]	5-10%	D-50	
521	[7%]	5-10%	D-51	
551	not detected	not detected	D-22	
561	not detected	not detected	D-15	
581	[1%]	under 5%	D-16	
591	12%	10-20%	D-50	
681	not detected	not detected	D-26	
682	not detected	not detected	D-26	

¹Prevalence means the proportion of sampled animals infected by CWD calculated as [number infected / total number sampled] and expressed here as a percentage (e.g., 10%). A prevalence of 20% is the same as a ratio of 1 in 5 (i.e. 1 in every 5 animals tested is infected), 10% is 1 in 10, 5% is 1 in 20, 1% is 1 in 100.

²CPW's best estimate of prevalence based on testing 100 or more submissions from this Game Management Unit (GMU). Values in brackets [] are based on the overall herd (Data Analysis Unit) prevalence estimate. Estimates shown for mule deer (MD) and white-tailed deer (WTD) where different. Data derived from the most recent year(s) when sample submissions were required in the GMU.

³Range of estimated prevalence within the corresponding herd/Data Analysis Unit (DAU) based on at least one year of mandatory sample submissions regardless of sample size. **Bold (5-10%)** or **bold red (10-20%; shaded >20%)** denotes GMUs/DAUs that exceed the 5% threshold that now compels CWD management action under CPW's response plan.

⁴"Not detected" means no cases reported from this DAU (herd) as of January 2023. **Does not assure absence of CWD.**

Estimated chronic wasting disease (CWD) prevalence¹ from harvested male deer. (See notes at right for further explanation.)

Game Management Unit (GMU)	GMU-level estimated prevalence ² (from mandatory submissions)	Prevalence Range ³	Herd/DAU
691	[<1%]	under 5%	D-34
711	3%	under 5%	D-24
741	not detected	not detected	D-52
751	[<1%]	under 5%	D-30
771	[<1%]	under 5%	D-30
791	not detected	not detected	D-36
851	[1%]	under 5%	D-32
861	[<1%]	under 5%	D-34
951	[26%]	over 20%	D-44

Notes

¹Prevalence means the proportion of sampled animals infected by CWD calculated as [number infected / total number sampled] and expressed here as a percentage (e.g., 10%). A prevalence of 20% is the same as a ratio of 1 in 5 (i.e. 1 in every 5 animals tested is infected), 10% is 1 in 10, 5% is 1 in 20, 1% is 1 in 100.

²CPW's best estimate of prevalence based on testing 100 or more submissions from this Game Management Unit (GMU). Values in brackets [] are based on the overall herd (Data Analysis Unit) prevalence estimate. Estimates shown for mule deer (MD) and white-tailed deer (WTD) where different. Data derived from the most recent year(s) when sample submissions were required in the GMU.

³Range of estimated prevalence within the corresponding herd/Data Analysis Unit (DAU) based on at least one year of mandatory sample submissions regardless of sample size. **Bold (5-10%)** or **bold red (10-20%;shaded >20%)** denotes GMUs/DAUs that exceed the 5% threshold that now compels CWD management action under CPW's response plan.

⁴"Not detected" means no cases reported from this DAU (herd) as of January 2023. **Does not assure absence of CWD.**